

# Continuous models of computation: computability, complexity, universality

Amaury Pouly

Joint work with Olivier Bournez and Daniel Graça

Université de Paris, IRIF, CNRS

14 june 2019



INSTITUT  
DE RECHERCHE  
EN INFORMATIQUE  
FONDAMENTALE



université  
PARIS  
DIDEROT  
Cité des Sciences



# What is a computer ?


# What is a computer?



# What is a computer?




VS



# Analog Computers



Differential Analyser  
“Mathematica of the 1920s”




Admiralty Fire Control Table  
British Navy ships (WW2)



# Church Thesis

## Complexity


discrete



## Effective Church Thesis

All **reasonable** models of computation are equivalent for complexity.

# Polynomial Differential Equations




General Purpose  
Analog Computer



Differential Analyzer



# Polynomial Differential Equations



General Purpose  
Analog Computer



Differential Analyzer

polynomial differential  
equations :

$$\begin{cases} y(0) = y_0 \\ y'(t) = p(y(t)) \end{cases}$$

# Polynomial Differential Equations

$$\boxed{k} \rightarrow k \quad \begin{matrix} u \\ v \end{matrix} \rightarrow \boxed{\times} \rightarrow uv$$

$$\begin{matrix} u \\ v \end{matrix} \rightarrow \boxed{+} \rightarrow u+v \quad u \rightarrow \boxed{\int} \rightarrow \int u$$

General Purpose  
Analog Computer



Differential Analyzer

Newton mechanics


Reaction networks :

- ▶ chemical
- ▶ enzymatic

polynomial differential  
equations :

$$\begin{cases} y(0) = y_0 \\ y'(t) = p(y(t)) \end{cases}$$

# Polynomial Differential Equations



General Purpose  
Analog Computer



Differential Analyzer

Newton mechanics

Reaction networks :


- ▶ chemical
- ▶ enzymatic

polynomial differential  
equations :

$$\begin{cases} y(0) = y_0 \\ y'(t) = p(y(t)) \end{cases}$$


- ▶ Rich class
- ▶ Stable (+,  $\times$ ,  $\circ$ ,  $/$ , ED)
- ▶ No closed-form solution

# Example of dynamical system



$$\ddot{\theta} + \frac{g}{\ell} \sin(\theta) = 0$$


# Example of dynamical system




$$\ddot{\theta} + \frac{g}{\ell} \sin(\theta) = 0$$

$$\begin{cases} y_1 = \theta \\ y_2 = \dot{\theta} \\ y_3 = \sin(\theta) \\ y_4 = \cos(\theta) \end{cases} \Leftrightarrow \begin{cases} y_1' = y_2 \\ y_2' = -\frac{g}{\ell} y_3 \\ y_3' = y_2 y_4 \\ y_4' = -y_2 y_3 \end{cases}$$

# Example of dynamical system




$$\ddot{\theta} + \frac{g}{\ell} \sin(\theta) = 0$$




$$\begin{cases} y_1 = \theta \\ y_2 = \dot{\theta} \\ y_3 = \sin(\theta) \\ y_4 = \cos(\theta) \end{cases} \Leftrightarrow \begin{cases} y_1' = y_2 \\ y_2' = -\frac{g}{\ell} y_3 \\ y_3' = y_2 y_4 \\ y_4' = -y_2 y_3 \end{cases}$$

# Example of dynamical system



$$\ddot{\theta} + \frac{g}{\ell} \sin(\theta) = 0$$



$$\begin{cases} y_1 = \theta \\ y_2 = \dot{\theta} \\ y_3 = \sin(\theta) \\ y_4 = \cos(\theta) \end{cases} \Leftrightarrow \begin{cases} y_1' = y_2 \\ y_2' = -\frac{g}{\ell} y_3 \\ y_3' = y_2 y_4 \\ y_4' = -y_2 y_3 \end{cases}$$

## Historical remark : the word “analog”


The pendulum and the circuit have the same equation. One can study one using the other by **analogy**.

# Computing with differential equations

## Generable functions

$$\begin{cases} y(0) = y_0 \\ y'(x) = p(y(x)) \end{cases} \quad x \in \mathbb{R}$$

$$f(x) = y_1(x)$$



Shannon's notion




# Computing with differential equations

## Generable functions

$$\begin{cases} y(0) = y_0 \\ y'(x) = p(y(x)) \end{cases} \quad x \in \mathbb{R}$$

$$f(x) = y_1(x)$$



Shannon's notion

sin, cos, exp, log, ...


Strictly weaker than Turing  
machines [Shannon, 1941]

# Computing with differential equations

## Generable functions

$$\begin{cases} y(0) = y_0 \\ y'(x) = p(y(x)) \end{cases} \quad x \in \mathbb{R}$$

$$f(x) = y_1(x)$$



Shannon's notion


sin, cos, exp, log, ...

Strictly weaker than Turing machines [Shannon, 1941]

## Computable

$$\begin{cases} y(0) = q(x) \\ y'(t) = p(y(t)) \end{cases} \quad \begin{matrix} x \in \mathbb{R} \\ t \in \mathbb{R}_+ \end{matrix}$$

$$f(x) = \lim_{t \rightarrow \infty} y_1(t)$$




Modern notion

# Computing with differential equations

## Generable functions

$$\begin{cases} y(0) = y_0 \\ y'(x) = p(y(x)) \end{cases} \quad x \in \mathbb{R}$$

$$f(x) = y_1(x)$$



Shannon's notion


sin, cos, exp, log, ...

Strictly weaker than Turing machines [Shannon, 1941]

## Computable

$$\begin{cases} y(0) = q(x) \\ y'(t) = p(y(t)) \end{cases} \quad \begin{matrix} x \in \mathbb{R} \\ t \in \mathbb{R}_+ \end{matrix}$$

$$f(x) = \lim_{t \rightarrow \infty} y_1(t)$$



Modern notion

sin, cos, exp, log,  $\Gamma$ ,  $\zeta$ , ...

Turing powerful  
[Bournez et al., 2007]

# From discrete to real computability

Computable Analysis : “Turing” computability over real numbers

# From discrete to real computability

Computable Analysis : “Turing” computability over real numbers

Definition (Ko, 1991 ; Weihrauch, 2000)

$x \in \mathbb{R}$  is computable iff  $\exists$  a computable  $f : \mathbb{N} \rightarrow \mathbb{Q}$  such that :

$$|x - f(n)| \leq 10^{-n} \quad n \in \mathbb{N}$$

Examples : rational numbers,  $\pi$ ,  $e$ , ...

<b>n</b>	<b>f(n)</b>	<b><math> \pi - f(n) </math></b>
0	3	$0.14 \leq 10^{-0}$
1	3.1	$0.04 \leq 10^{-1}$
2	3.14	$0.001 \leq 10^{-2}$
10	3.1415926535	$0.9 \cdot 10^{-10} \leq 10^{-10}$

# From discrete to real computability

Computable Analysis : “Turing” computability over real numbers

Definition (Ko, 1991 ; Weihrauch, 2000)

$x \in \mathbb{R}$  is computable iff  $\exists$  a computable  $f : \mathbb{N} \rightarrow \mathbb{Q}$  such that :

$$|x - f(n)| \leq 10^{-n} \quad n \in \mathbb{N}$$


Examples : rational numbers,  $\pi$ ,  $e$ , ...

<b>n</b>	<b>f(n)</b>	<b><math> \pi - f(n) </math></b>
0	3	$0.14 \leq 10^{-0}$
1	3.1	$0.04 \leq 10^{-1}$
2	3.14	$0.001 \leq 10^{-2}$
10	3.1415926535	$0.9 \cdot 10^{-10} \leq 10^{-10}$


Beware : there exists uncomputable real numbers !

$$x = \sum_{n \in \Gamma} 2^{-n}, \quad \Gamma = \{n : \text{the } n^{\text{th}} \text{ Turing machine halts}\}$$

# From discrete to real computability



# From discrete to real computability




## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}$ ,  
computable functions such that :

- ▶ effective approx over  $\mathbb{Q} : |f(r) - \psi(r, n)| \leq 10^{-n}$



# From discrete to real computability




## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}$ ,  
computable functions such that :

- ▶ effective approx over  $\mathbb{Q} : |f(r) - \psi(r, n)| \leq 10^{-n}$

# From discrete to real computability




## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}$ ,  
computable functions such that :

- ▶ effective approx over  $\mathbb{Q} : |f(r) - \psi(r, n)| \leq 10^{-n}$

# From discrete to real computability




## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

# From discrete to real computability




## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

# From discrete to real computability



## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

# From discrete to real computability

## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

All computable functions are continuous !

Examples : polynomials,  $\sin$ ,  $\exp$ ,  $\sqrt{\cdot}$

Beware : there exists (continuous) uncomputable real functions !

# From discrete to real computability

## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

All computable functions are continuous !

Examples : polynomials,  $\sin$ ,  $\exp$ ,  $\sqrt{\cdot}$

Beware : there exists (continuous) uncomputable real functions !

## Polytime complexity

Add “polynomial time computable” everywhere.

# From discrete to real computability

## Definition (Computable function)

$f : [a, b] \rightarrow \mathbb{R}$  is computable iff  $\exists m : \mathbb{N} \rightarrow \mathbb{N}, \psi : \mathbb{Q} \times \mathbb{N} \rightarrow \mathbb{Q}$  computable functions such that :

- ▶ effective approx over  $\mathbb{Q}$  :  $|f(r) - \psi(r, n)| \leq 10^{-n}$
- ▶ effective continuity :  $|x - y| \leq 10^{-m(n)} \Rightarrow |f(x) - f(y)| \leq 10^{-n}$   
 $m$  : modulus of continuity

All computable functions are continuous !

Examples : polynomials,  $\sin$ ,  $\exp$ ,  $\sqrt{\cdot}$ .

Beware : there exists (continuous) uncomputable real functions !

## Polytime complexity

Add “polynomial time computable” everywhere.

Remark : there are other theories of computability over  $\mathbb{R}$ , notably BSS (Blum-Shub-Smale).




# Equivalence with computable analysis

## Definition (Bournez et al, 2007)

$f$  **computable by GPAC** if  $\exists p$  polynomial such that  $\forall x \in [a, b]$

$$y(0) = (x, 0, \dots, 0) \quad y'(t) = p(y(t))$$

satisfies  $|f(x) - y_1(t)| \leq y_2(t)$  et  $y_2(t) \xrightarrow[t \rightarrow \infty]{} 0$ .



$$y_1(t) \xrightarrow[t \rightarrow \infty]{} f(x)$$

$$y_2(t) = \text{error bound}$$


# Equivalence with computable analysis

## Definition (Bournez et al, 2007)

$f$  **computable by GPAC** if  $\exists p$  polynomial such that  $\forall x \in [a, b]$

$$y(0) = (x, 0, \dots, 0) \quad y'(t) = p(y(t))$$

satisfies  $|f(x) - y_1(t)| \leq y_2(t)$  et  $y_2(t) \xrightarrow[t \rightarrow \infty]{} 0$ .



$$y_1(t) \xrightarrow[t \rightarrow \infty]{} f(x)$$

$$y_2(t) = \text{error bound}$$

## Theorem (Bournez et al, 2007)

$f : [a, b] \rightarrow \mathbb{R}$  *computable*  $\Leftrightarrow f$  *computable by GPAC*

# Complexity of analog systems


- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$

# Complexity of analog systems

- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$
- ▶ GPAC :

## Tentative definition

$$T(x) = ??$$




# Complexity of analog systems

- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$
- ▶ GPAC :

## Tentative definition

$$T(x, \mu) =$$




# Complexity of analog systems

- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$
- ▶ GPAC :

## Tentative definition

$T(x, \mu) =$  first time  $t$  so that  $|y_1(t) - f(x)| \leq e^{-\mu}$




# Complexity of analog systems


- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$
- ▶ GPAC :

## Tentative definition

$T(x, \mu) =$  first time  $t$  so that  $|y_1(t) - f(x)| \leq e^{-\mu}$



$\leadsto$




# Complexity of analog systems

- ▶ Turing machines :  $T(x)$  = number of steps to compute on  $x$
- ▶ GPAC :

## Tentative definition


$T(x, \mu) = \text{first time } t \text{ so that } |y_1(t) - f(x)| \leq e^{-\mu}$

$$y(0) = (x, 0, \dots, 0) \quad y' = p(y)$$




$\leadsto$

$$z(t) = y(e^t)$$



$$w(t) = y(e^{e^t})$$






# Complexity of analog systems

- ▶ **Turing machines** :  $T(x)$  = number of steps to compute on  $x$
- ▶ **GPAC** : time contraction problem → **open problem**

## Tentative definition


$T(x, \mu) =$  first time  $t$  so that  $|y_1(t) - f(x)| \leq e^{-\mu}$

$$y(0) = (x, 0, \dots, 0) \quad y' = p(y)$$



$\leadsto$


$$z(t) = y(e^t)$$




Something is wrong...

All functions have constant time complexity.


$$w(t) = y(e^{e^t})$$




# Time-space correlation of the GPAC




$\leadsto$




# Time-space correlation of the GPAC




$\leadsto$




extra component :  $w(t) = e^t$



# Time-space correlation of the GPAC



$\rightsquigarrow$




## Observation

Time scaling costs “space”.

$\rightsquigarrow$

Time complexity for the GPAC  
must involve time and **space** !


extra component :  $w(t) = e^t$



# Characterization of polynomial time

**Definition :**  $\mathcal{L} \in \text{ANALOG-PTIME} \Leftrightarrow \exists p \text{ polynomial, } \forall \text{ word } w$


$$y(0) = (\psi(w), |w|, 0, \dots, 0) \quad y' = p(y) \quad \psi(w) = \sum_{i=1}^{|w|} w_i 2^{-i}$$



# Characterization of polynomial time

**Definition :**  $\mathcal{L} \in \text{ANALOG-PTIME} \Leftrightarrow \exists p \text{ polynomial, } \forall \text{ word } w$

$$y(0) = (\psi(w), |w|, 0, \dots, 0) \quad y' = p(y) \quad \psi(w) = \sum_{i=1}^{|w|} w_i 2^{-i}$$




satisfies

1. if  $y_1(t) \geq 1$  then  $w \in \mathcal{L}$

# Characterization of polynomial time

**Definition :**  $\mathcal{L} \in \text{ANALOG-PTIME} \Leftrightarrow \exists p \text{ polynomial, } \forall \text{ word } w$

$$y(0) = (\psi(w), |w|, 0, \dots, 0) \quad y' = p(y) \quad \psi(w) = \sum_{i=1}^{|w|} w_i 2^{-i}$$




satisfies

2. if  $y_1(t) \leq -1$  then  $w \notin \mathcal{L}$

# Characterization of polynomial time

**Definition :**  $\mathcal{L} \in \text{ANALOG-PTIME} \Leftrightarrow \exists p \text{ polynomial, } \forall \text{ word } w$

$$y(0) = (\psi(w), |w|, 0, \dots, 0) \quad y' = p(y) \quad \psi(w) = \sum_{i=1}^{|w|} w_i 2^{-i}$$



satisfies


3. if  $\ell(t) \geq \text{poly}(|w|)$  then  $|y_1(t)| \geq 1$



# Characterization of polynomial time

**Definition :**  $\mathcal{L} \in \text{ANALOG-PTIME} \Leftrightarrow \exists p \text{ polynomial}, \forall \text{ word } w$

$$y(0) = (\psi(w), |w|, 0, \dots, 0) \quad y' = p(y) \quad \psi(w) = \sum_{i=1}^{|w|} w_i 2^{-i}$$




**Theorem**


$$\text{PTIME} = \text{ANALOG-PTIME}$$

# Summary

## ANALOG-PTIME



## ANALOG- $P_{\mathbb{R}}$




## Theorem

- ▶  $\mathcal{L} \in \text{PTIME}$  if and only if  $\mathcal{L} \in \text{ANALOG-PTIME}$
- ▶  $f : [a, b] \rightarrow \mathbb{R}$  computable in polynomial time  $\Leftrightarrow f \in \text{ANALOG-}P_{\mathbb{R}}$


- ▶ Analog complexity theory based on **length**
- ▶ Time of Turing machine  $\Leftrightarrow$  length of the GPAC
- ▶ Purely continuous characterization of PTIME

# Summary

## ANALOG-PTIME



## ANALOG- $P_{\mathbb{R}}$



## Theorem

- ▶  $\mathcal{L} \in \text{PTIME}$  if and only if  $\mathcal{L} \in \text{ANALOG-PTIME}$
- ▶  $f : [a, b] \rightarrow \mathbb{R}$  computable in polynomial time  $\Leftrightarrow f \in \text{ANALOG-}P_{\mathbb{R}}$

- ▶ Analog complexity theory based on **length**
- ▶ Time of Turing machine  $\Leftrightarrow$  length of the GPAC
- ▶ Purely continuous characterization of PTIME
- ▶ Only **rational coefficients** needed

# In the remaining time...

Two applications of the techniques we have developed :

~> Chemical Reaction Networks


Universal differential equation

# Chemical Reaction Networks

**Definition :** a **reaction system** is a finite set of

- ▶ molecular species  $y_1, \dots, y_n$
- ▶ reactions of the form  $\sum_i a_i y_i \xrightarrow{f} \sum_i b_i y_i$  ( $a_i, b_i \in \mathbb{N}$ ,  $f = \text{rate}$ )

Example (any resemblance to chemistry is purely coincidental) :




# Chemical Reaction Networks

**Definition :** a **reaction system** is a finite set of

- ▶ molecular species  $y_1, \dots, y_n$
- ▶ reactions of the form  $\sum_i a_i y_i \xrightarrow{f} \sum_i b_i y_i \quad (a_i, b_i \in \mathbb{N}, f = \text{rate})$

Example (any resemblance to chemistry is purely coincidental) :



**Assumption :** law of mass action


$$\sum_i a_i y_i \xrightarrow{k} \sum_i b_i y_i \rightsquigarrow f(y) = k \prod_i y_i^{a_i}$$

# Chemical Reaction Networks

**Definition :** a **reaction system** is a finite set of

- ▶ molecular species  $y_1, \dots, y_n$
- ▶ reactions of the form  $\sum_i a_i y_i \xrightarrow{f} \sum_i b_i y_i$  ( $a_i, b_i \in \mathbb{N}$ ,  $f = \text{rate}$ )

Example (any resemblance to chemistry is purely coincidental) :



**Assumption :** law of mass action

$$\sum_i a_i y_i \xrightarrow{k} \sum_i b_i y_i \rightsquigarrow f(y) = k \prod_i y_i^{a_i}$$

**Semantics :**


- ▶ discrete
- ▶ differential
- ▶ stochastic

# Chemical Reaction Networks

**Definition :** a **reaction system** is a finite set of

- ▶ molecular species  $y_1, \dots, y_n$
- ▶ reactions of the form  $\sum_i a_i y_i \xrightarrow{f} \sum_i b_i y_i \quad (a_i, b_i \in \mathbb{N}, f = \text{rate})$

**Example** (any resemblance to chemistry is purely coincidental) :



**Assumption :** law of mass action

$$\sum_i a_i y_i \xrightarrow{k} \sum_i b_i y_i \rightsquigarrow f(y) = k \prod_i y_i^{a_i}$$

**Semantics :**

- ▶ discrete
- ▶ differential  $\rightarrow$
- ▶ stochastic

$$y'_i = \sum_{\text{reaction } R} (b_i^R - a_i^R) f^R(y)$$




# Chemical Reaction Networks

**Definition :** a **reaction system** is a finite set of

- ▶ molecular species  $y_1, \dots, y_n$
- ▶ reactions of the form  $\sum_i a_i y_i \xrightarrow{f} \sum_i b_i y_i \quad (a_i, b_i \in \mathbb{N}, f = \text{rate})$

**Example** (any resemblance to chemistry is purely coincidental) :



**Assumption :** law of mass action

$$\sum_i a_i y_i \xrightarrow{k} \sum_i b_i y_i \rightsquigarrow f(y) = k \prod_i y_i^{a_i}$$

**Semantics :**

- ▶ discrete
- ▶ differential  $\rightarrow$
- ▶ stochastic

$$y_i' = \sum_{\text{reaction } R} (b_i^R - a_i^R) k^R \prod_j y_j^{a_j^R}$$

# Chemical Reaction Networks (CRNs)

- ▶ CRNs with differential semantics and mass action law = polynomial ODEs
- ▶ polynomial ODEs are Turing complete

# Chemical Reaction Networks (CRNs)

- ▶ CRNs with differential semantics and mass action law = polynomial ODEs
- ▶ polynomial ODEs are Turing complete

CRNs are Turing complete ?

# Chemical Reaction Networks (CRNs)

CRNs are Turing complete ? Two “slight” problems :

- ▶ concentrations cannot be negative ( $y_i < 0$ )
- ▶ arbitrary reactions are not realistic

# Chemical Reaction Networks (CRNs)

CRNs are Turing complete ? Two “slight” problems :

- ▶ concentrations cannot be negative ( $y_i < 0$ )
- ▶ arbitrary reactions are not realistic
- ▶ easy to solve
- ▶ what is realistic ?

# Chemical Reaction Networks (CRNs)

CRNs are Turing complete ? Two “slight” problems :

- ▶ concentrations cannot be negative ( $y_i < 0$ )
- ▶ arbitrary reactions are not realistic
- ▶ easy to solve
- ▶ what is realistic ?

**Definition :** a reaction is **elementary** if it has at most two reactants  
⇒ can be implemented with DNA, RNA or proteins

# Chemical Reaction Networks (CRNs)

CRNs are Turing complete? Two “slight” problems :

- ▶ concentrations cannot be negative ( $y_i < 0$ )
- ▶ arbitrary reactions are not realistic
- ▶ easy to solve
- ▶ what is realistic?

**Definition :** a reaction is **elementary** if it has at most two reactants  
⇒ can be implemented with DNA, RNA or proteins

Elementary reactions correspond to **quadratic** ODEs :



# Chemical Reaction Networks (CRNs)

CRNs are Turing complete? Two “slight” problems :

- ▶ concentrations cannot be negative ( $y_i < 0$ )
- ▶ arbitrary reactions are not realistic
- ▶ easy to solve
- ▶ what is realistic?

**Definition :** a reaction is **elementary** if it has at most two reactants  
⇒ can be implemented with DNA, RNA or proteins

Elementary reactions correspond to **quadratic** ODEs :



## Theorem (Folklore)

*Every polynomial ODE can be rewritten as a quadratic ODE.*



# Chemical Reaction Networks (CRNs)

**Definition :** a reaction is **elementary** if it has at most two reactants  
⇒ can be implemented with DNA, RNA or proteins

Elementary reactions correspond to **quadratic** ODEs :


$$ay + bz \xrightarrow{k} \dots \quad \rightsquigarrow \quad f(y, z) = ky^a z^b$$

**Theorem (Work with François Fages, Guillaume Le Guludec)**

*Elementary mass-action-law reaction system on finite universes of molecules are Turing-complete under the differential semantics.*

Notes :

- ▶ proof preserves polynomial length
- ▶ in fact the following elementary reactions suffice :



# In the remaining time...


Two applications of the techniques we have developed :

Chemical Reaction Networks

~> Universal differential equation


# Universal differential equations

## Generable functions



subclass of analytic functions


## Computable functions




any computable function

# Universal differential equations

## Generable functions




## Computable functions




subclass of analytic functions

any computable function



# Universal differential algebraic equation (DAE)



## Theorem (Rubel, 1981)


For any continuous functions  $f$  and  $\varepsilon$ , there exists  $y : \mathbb{R} \rightarrow \mathbb{R}$  solution to

$$\begin{aligned} &3y'^4 y'' y''''^2 - 4y'^4 y'''^2 y'''' + 6y'^3 y''^2 y''' y'''' + 24y'^2 y''^4 y'''' \\ &- 12y'^3 y'' y'''^3 - 29y'^2 y''^3 y'''^2 + 12y''^7 = 0 \end{aligned}$$

such that  $\forall t \in \mathbb{R}$ ,

$$|y(t) - f(t)| \leq \varepsilon(t).$$

# Universal differential algebraic equation (DAE)



## Theorem (Rubel, 1981)


*There exists a **fixed** polynomial  $p$  and  $k \in \mathbb{N}$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists a solution  $y : \mathbb{R} \rightarrow \mathbb{R}$  to*

$$p(y, y', \dots, y^{(k)}) = 0$$

*such that  $\forall t \in \mathbb{R}$ ,*

$$|y(t) - f(t)| \leq \varepsilon(t).$$

# Universal differential algebraic equation (DAE)



## Theorem (Rubel, 1981)

There exists a **fixed** polynomial  $p$  and  $k \in \mathbb{N}$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists a solution  $y : \mathbb{R} \rightarrow \mathbb{R}$  to

$$p(y, y', \dots, y^{(k)}) = 0$$

such that  $\forall t \in \mathbb{R}$ ,

$$|y(t) - f(t)| \leq \varepsilon(t).$$

**Problem** : this is «weak» result.

# The problem with Rubel's DAE

The solution  $y$  is not unique, **even with added initial conditions** :

$$p(y, y', \dots, y^{(k)}) = 0, \quad y(0) = \alpha_0, y'(0) = \alpha_1, \dots, y^{(k)}(0) = \alpha_k$$

In fact, this is fundamental for Rubel's proof to work !



# The problem with Rubel's DAE

The solution  $y$  is not unique, **even with added initial conditions** :

$$p(y, y', \dots, y^{(k)}) = 0, \quad y(0) = \alpha_0, y'(0) = \alpha_1, \dots, y^{(k)}(0) = \alpha_k$$

In fact, this is fundamental for Rubel's proof to work !

- ▶ Rubel's statement : this DAE is universal
- ▶ More realistic interpretation : this DAE allows almost anything

## Open Problem (Rubel, 1981)


Is there a universal ODE  $y' = p(y)$  ?

**Note** : explicit polynomial ODE  $\Rightarrow$  unique solution

# Rubel's proof in one slide

- Take  $f(t) = e^{\frac{-1}{1-t^2}}$  for  $-1 < t < 1$  and  $f(t) = 0$  otherwise.

It satisfies  $(1 - t^2)^2 f''(t) + 2tf'(t) = 0$ .



# Rubel's proof in one slide


- Take  $f(t) = e^{\frac{-1}{1-t^2}}$  for  $-1 < t < 1$  and  $f(t) = 0$  otherwise.

It satisfies  $(1 - t^2)^2 f''(t) + 2t f'(t) = 0$ .

- For any  $a, b, c \in \mathbb{R}$ ,  $y(t) = cf(at + b)$  satisfies

$$\begin{aligned} 3y'^4 y'' y''''^2 & - 4y'^4 y''^2 y'''' + 6y'^3 y''^2 y''' y'''' + 24y'^2 y''^4 y'''' \\ & - 12y'^3 y'' y''''^3 - 29y'^2 y''^3 y''''^2 + 12y''^7 = 0 \end{aligned}$$

Translation and rescaling :



# Rubel's proof in one slide


- ▶ Take  $f(t) = e^{\frac{-1}{1-t^2}}$  for  $-1 < t < 1$  and  $f(t) = 0$  otherwise.

It satisfies  $(1 - t^2)^2 f''(t) + 2t f'(t) = 0$ .

- ▶ For any  $a, b, c \in \mathbb{R}$ ,  $y(t) = cf(at + b)$  satisfies

$$3y'^4 y'' y''''^2 - 4y'^4 y''^2 y'''' + 6y'^3 y''^2 y''' y'''' + 24y'^2 y''^4 y'''' - 12y'^3 y'' y''''^3 - 29y'^2 y''^3 y''''^2 + 12y''^7 = 0$$

- ▶ Can glue together arbitrary many such pieces



# Rubel's proof in one slide


- Take  $f(t) = e^{\frac{-1}{1-t^2}}$  for  $-1 < t < 1$  and  $f(t) = 0$  otherwise.

It satisfies  $(1 - t^2)^2 f''(t) + 2t f'(t) = 0$ .

- For any  $a, b, c \in \mathbb{R}$ ,  $y(t) = cf(at + b)$  satisfies

$$3y'^4 y'' y''''^2 - 4y'^4 y''^2 y'''' + 6y'^3 y''^2 y''' y'''' + 24y'^2 y''^4 y'''' - 12y'^3 y'' y''''^3 - 29y'^2 y''^3 y''''^2 + 12y''^7 = 0$$

- Can glue together arbitrary many such pieces
- Can arrange so that  $\int f$  is solution : **piecewise pseudo-linear**



# Rubel's proof in one slide


- Take  $f(t) = e^{\frac{-1}{1-t^2}}$  for  $-1 < t < 1$  and  $f(t) = 0$  otherwise.

It satisfies  $(1 - t^2)^2 f''(t) + 2tf'(t) = 0$ .

- For any  $a, b, c \in \mathbb{R}$ ,  $y(t) = cf(at + b)$  satisfies


$$3y'^4 y'' y''''^2 - 4y'^4 y''^2 y'''' + 6y'^3 y''^2 y''' y'''' + 24y'^2 y''^4 y'''' - 12y'^3 y'' y''''^3 - 29y'^2 y''^3 y''''^2 + 12y''^7 = 0$$

- Can glue together arbitrary many such pieces
- Can arrange so that  $\int f$  is solution : **piecewise pseudo-linear**



**Conclusion** : Rubel's equation allows any piecewise pseudo-linear functions, and those are **dense** in  $C^0$

# Universal initial value problem (IVP)



## Theorem


There exists a **fixed** (vector of) polynomial  $p$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists  $\alpha \in \mathbb{R}^d$  such that

$$y(0) = \alpha, \quad y'(t) = p(y(t))$$

has a **unique solution**  $y : \mathbb{R} \rightarrow \mathbb{R}^d$  and  $\forall t \in \mathbb{R}$ ,

$$|y_1(t) - f(t)| \leq \varepsilon(t).$$

# Universal initial value problem (IVP)



Notes :

- ▶ **system** of ODEs,
- ▶  $y$  is analytic,
- ▶ we need  $d \approx 300$ .

## Theorem

There exists a **fixed** (vector of) polynomial  $p$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists  $\alpha \in \mathbb{R}^d$  such that


$$y(0) = \alpha, \quad y'(t) = p(y(t))$$

has a **unique solution**  $y : \mathbb{R} \rightarrow \mathbb{R}^d$  and  $\forall t \in \mathbb{R}$ ,

$$|y_1(t) - f(t)| \leq \varepsilon(t).$$



# Universal initial value problem (IVP)



Notes :

- ▶ **system** of ODEs,
- ▶  $y$  is analytic,
- ▶ we need  $d \approx 300$ .

## Theorem

There exists a **fixed** (vector of) polynomial  $p$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists  $\alpha \in \mathbb{R}^d$  such that

$$y(0) = \alpha, \quad y'(t) = p(y(t))$$

has a **unique solution**  $y : \mathbb{R} \rightarrow \mathbb{R}^d$  and  $\forall t \in \mathbb{R}$ ,

$$|y_1(t) - f(t)| \leq \varepsilon(t).$$

**Remark :**  $\alpha$  is usually transcendental, but computable from  $f$  and  $\varepsilon$



Reaction networks :

- ▶ chemical
- ▶ enzymatic

$$y' = p(y)$$

?

$$y' = p(y) + e(t)$$

- ▶ Finer time complexity (linear)
- ▶ Nondeterminism
- ▶ Robustness
- ▶ « Space » complexity
- ▶ Other models
- ▶ Stochastic

# Backup slides

# Complexity of solving polynomial ODEs

$$y(0) = x \quad y'(t) = p(y(t))$$



# Complexity of solving polynomial ODEs

$$y(0) = x \quad y'(t) = p(y(t))$$

## Theorem

If  $y(t)$  exists, one can compute  $p, q$  such that  $\left| \frac{p}{q} - y(t) \right| \leq 2^{-n}$  in time  
 $\text{poly}(\text{size of } x \text{ and } p, n, \ell(t))$

where  $\ell(t) \approx$  length of the curve (between  $x$  and  $y(t)$ )




length of the curve = complexity = resource

# Characterization of real polynomial time

**Definition :**  $f : [a, b] \rightarrow \mathbb{R}$  in  $\text{ANALOG-P}_{\mathbb{R}} \Leftrightarrow \exists p$  polynomial,  $\forall x \in [a, b]$

$$y(0) = (x, 0, \dots, 0) \quad y' = p(y)$$



# Characterization of real polynomial time

**Definition :**  $f : [a, b] \rightarrow \mathbb{R}$  in  $\text{ANALOG-P}_{\mathbb{R}} \Leftrightarrow \exists p$  polynomial,  $\forall x \in [a, b]$

$$y(0) = (x, 0, \dots, 0) \quad y' = p(y)$$


satisfies :

1.  $|y_1(t) - f(x)| \leq 2^{-\ell(t)}$

«greater length  $\Rightarrow$  greater precision»

2.  $\ell(t) \geq t$

«length increases with time»



# Characterization of real polynomial time

**Definition :**  $f : [a, b] \rightarrow \mathbb{R}$  in  $\text{ANALOG-P}_{\mathbb{R}} \Leftrightarrow \exists p$  polynomial,  $\forall x \in [a, b]$

$$y(0) = (x, 0, \dots, 0) \quad y' = p(y)$$


satisfies :

1.  $|y_1(t) - f(x)| \leq 2^{-\ell(t)}$

«greater length  $\Rightarrow$  greater precision»

2.  $\ell(t) \geq t$

«length increases with time»




## Theorem

$f : [a, b] \rightarrow \mathbb{R}$  computable in polynomial time  $\Leftrightarrow f \in \text{ANALOG-P}_{\mathbb{R}}$ .



# Universal DAE revisited



## Theorem

There exists a **fixed** polynomial  $p$  and  $k \in \mathbb{N}$  such that for any continuous functions  $f$  and  $\varepsilon$ , there exists  $\alpha_0, \dots, \alpha_k \in \mathbb{R}$  such that

$$p(y, y', \dots, y^{(k)}) = 0, \quad y(0) = \alpha_0, y'(0) = \alpha_1, \dots, y^{(k)}(0) = \alpha_k$$

has a **unique analytic solution** and this solution satisfies such that

$$|y(t) - f(t)| \leq \varepsilon(t).$$